arXiv:1503.08418v1 [math.NT] 29 Mar 2015

A NOTE ON DEGENERATE POLY-BERNOULLI NUMBERS AND POLYNOMIALS

DAE SAN KIM AND TAEKYUN KIM

Abstract. In this paper, we consider the degenerate poly-Bernoulli polynomials and present new and explicit formulas for computing them in terms of the degenerate Bernoulli polynomials and Stirling numbers of the second kind.

1. Introduction

For $\lambda \in \mathbb{C}$, L. Carlitz considered the degenerate Bernoulli polynomials given by the generating function

$$\frac{t}{(1+\lambda t)^{\frac{1}{\lambda}}-1}(1+\lambda t)^{\frac{x}{\lambda}}=\sum_{n=0}^{\infty}\beta_n(x\mid\lambda)\frac{t^n}{n!},\quad(\text{see }[6,\,15,\,19]).\tag{1.1}$$

When $x=0$, $\beta_n(\lambda)=\beta_n(0\mid\lambda)$ are called the degenerate Bernoulli numbers.

Thus, by (1.1), we get

$$\beta_n(x\mid\lambda)=\sum_{l=0}^{n}\binom{n}{l}\beta_l(\lambda)(x\mid\lambda)_{n-l},\tag{1.2}$$

where $(x\mid\lambda)_n=x(x-\lambda)(x-2\lambda)\cdots(x-\lambda(n-1))$.

The classical polylograithm function Li_k is

$$\mathrm{Li}_k(x)=\sum_{n=1}^{\infty}\frac{x^n}{n^k},\quad(k\in\mathbb{Z}),\quad(\text{see }[10\text{–}17,\,20]).\tag{1.3}$$

From (1.1), we note that

$$\begin{aligned}&\sum_{n=0}^{\infty}\lim_{\lambda\to0}\beta_n(x\mid\lambda)\frac{t^n}{n!}\\&=\lim_{\lambda\to0}\frac{t}{(1+\lambda t)^{\frac{1}{\lambda}}-1}(1+\lambda t)^{\frac{x}{\lambda}}\\&=\frac{t}{e^t-1}e^{xt}\\&=\sum_{n=0}^{\infty}B_n(x)\frac{t^n}{n!},\end{aligned}\tag{1.4}$$

where $B_n(x)$ are called the Bernoulli polynomials (see [1–27]).

2010 *Mathematics Subject Classification.* 11B68, 11B73, 11B83.

Key words and phrases. degenerate poly-Bernoulli polynomial, degenerate Bernoulli polynomial, Stirling number of the second kind .

Thus, by (1.4), we get

$$\lim_{\lambda\to 0}\beta_n\left(x\mid\lambda\right)=B_n\left(x\right),\quad (n\geq 0)\,. \tag{1.5}$$

In [4, 10], the poly-Bernoulli polynomials are given by

$$\frac{\mathrm{Li}_k\left(1-e^{-t}\right)}{e^t-1}e^{xt}=\sum_{n=0}^{\infty}B_n^{(k)}\left(x\right)\frac{t^n}{n!}. \tag{1.6}$$

For $k=1$, we have

$$\frac{\mathrm{Li}_1\left(1-e^{-t}\right)}{e^t-1}e^{xt}=\frac{t}{e^t-1}e^{xt}=\sum_{n=0}^{\infty}B_n\left(x\right)\frac{t^n}{n!}. \tag{1.7}$$

By (1.4) and (1.7), we get $B_n^{(1)}\left(x\right)=B_n\left(x\right)$.
The Stirling numbers of the second kind are given by

$$x^n=\sum_{l=0}^{n}S_2\left(n,l\right)\left(x\right)_l\,,\quad \text{(see [1–27])}\,. \tag{1.8}$$

and the Stirling numbers of the first kind are defined by

$$\left(x\right)_n=x\left(x-1\right)\cdots\left(x-n+1\right)=\sum_{l=0}^{n}S_1\left(n,l\right)x^l,\quad (n\geq 0)\,. \tag{1.9}$$

The purpose of this paper is to construct the degenerate poly-Bernoulli polynomials and present new and explicit formulas for computing them in terms of the degenerate Bernoulli polynomials and Stirling numbers of the second kind.

2. DEGENERATE POLY-BERNOULLI NUMBERS AND POLYNOMIALS

For $\lambda\in\mathbb{C}$, $k\in\mathbb{Z}$, we consider the degenerate poly-Bernoulli polynomials given by the generating function

$$\frac{\mathrm{Li}_k\left(1-e^{-t}\right)}{\left(1+\lambda t\right)^{\frac{1}{\lambda}}-1}\left(1+\lambda t\right)^{\frac{x}{\lambda}}=\sum_{n=0}^{\infty}\beta_n^{(k)}\left(x\mid\lambda\right)\frac{t^n}{n!}. \tag{2.1}$$

When $x=0$, $\beta_n^{(k)}\left(\lambda\right)=\beta_n^{(k)}\left(0\mid\lambda\right)$ are called the degenerate poly-Bernoulli numbers. Note that $\beta_n^{(1)}\left(x\mid\lambda\right)=\beta_n\left(x\mid\lambda\right)$ and $\lim_{\lambda\to 0}\beta_n^{(k)}\left(x\mid\lambda\right)=B_n^{(k)}\left(x\right)$.
From (2.1), we can derive the following equation:

$$\begin{aligned}\sum_{n=0}^{\infty}\beta_n^{(k)}\left(x\mid\lambda\right)\frac{t^n}{n!}&=\left(\frac{\mathrm{Li}_k\left(1-e^{-t}\right)}{\left(1+\lambda t\right)^{\frac{1}{\lambda}}-1}\right)\left(1+\lambda t\right)^{\frac{x}{\lambda}}\\&=\left(\sum_{l=0}^{\infty}\beta_l^{(k)}\left(\lambda\right)\frac{t^l}{l!}\right)\left(\sum_{m=0}^{\infty}\left(x\mid\lambda\right)_m\frac{t^m}{m!}\right)\\&=\sum_{n=0}^{\infty}\left(\sum_{l=0}^{n}\binom{n}{l}\beta_l^{(k)}\left(\lambda\right)\left(x\mid\lambda\right)_{n-l}\right)\frac{t^n}{n!}.\end{aligned} \tag{2.2}$$

Thus, by (2.2), we get

$$\beta_n^{(k)}\left(x\mid\lambda\right)=\sum_{l=0}^{n}\binom{n}{l}\beta_l^{(k)}\left(\lambda\right)\left(x\mid\lambda\right)_{n-l}\,. \tag{2.3}$$

Now, we observe that

$$\begin{aligned}
&\frac{\mathrm{Li}_k\left(1-e^{-t}\right)}{(1+\lambda t)^{\frac{1}{\lambda}}-1}(1+t)^{\frac{x}{\lambda}} \\
&=\sum_{n=0}^{\infty}\beta_n^{(k)}(x \mid \lambda)\frac{t^n}{n!} \\
&=\frac{(1+t)^{\frac{x}{\lambda}}}{(1+\lambda t)^{\frac{1}{\lambda}}-1}\int_0^t \underbrace{\frac{1}{e^y-1}\int_0^y \frac{1}{e^y-1}\int_0^y \cdots \frac{1}{e^y-1}\int_0^y}_{(k-2)\mathrm{times}} \frac{y}{e^y-1}dy\cdots dy.
\end{aligned} \tag{2.4}$$

From (2.4), we have

$$\begin{aligned}
&\sum_{n=0}^{\infty}\beta_n^{(2)}(x \mid \lambda)\frac{t^n}{n!} \\
&=\frac{(1+t)^{\frac{x}{\lambda}}}{(1+\lambda t)^{\frac{1}{\lambda}}-1}\int_0^t \frac{y}{e^y-1}dy \\
&=\frac{(1+t)^{\frac{x}{\lambda}}}{(1+\lambda t)^{\frac{1}{\lambda}}-1}\sum_{l=0}^{\infty}\frac{B_l}{l!}\int_0^t y^l dy \\
&=\left(\frac{t}{(1+\lambda t)^{\frac{1}{\lambda}}-1}(1+\lambda t)^{\frac{x}{\lambda}}\right)\left(\sum_{l=0}^{\infty}\frac{B_l}{l+1}\frac{t^l}{l!}\right) \\
&=\sum_{n=0}^{\infty}\left\{\sum_{l=0}^{n}\binom{n}{l}\frac{B_l}{l+1}\beta_{n-l}(x \mid \lambda)\right\}\frac{t^n}{n!},
\end{aligned} \tag{2.5}$$

where $B_n = B_n(0)$ are Bernoulli numbers.

By comparing the coefficients on both sides of (2.5), we obtain the following theorem.

Theorem 2.1. *For $n \geq 0$, we have*

$$\begin{aligned}
\beta_n^{(2)}(x \mid \lambda) &= \sum_{l=0}^{n}\binom{n}{l}\frac{B_l}{l+1}\beta_{n-l}(x \mid \lambda) \\
&= \beta_n(x \mid \lambda) - \frac{n}{4}\beta_{n-1}(x \mid \lambda) + \sum_{l=2}^{n}\binom{n}{l}\frac{B_l}{l+1}\beta_{n-l}(x \mid \lambda).
\end{aligned}$$

Moreover,

$$\beta_n^{(k)}(x \mid \lambda) = \sum_{l=0}^{n}\binom{n}{l}\beta_l^{(k)}(\lambda)(x \mid \lambda)_{n-l}.$$

By (2.4), we easily get

$$\begin{aligned}
&\sum_{n=0}^{\infty}\beta_n^{(k)}(x \mid \lambda)\frac{t^n}{n!} \\
&=\frac{\mathrm{Li}_k\left(1-e^{-t}\right)}{(1+\lambda t)^{\frac{1}{\lambda}}-1}(1+t)^{\frac{x}{\lambda}}
\end{aligned} \tag{2.6}$$

$$= \frac{t}{(1+\lambda t)^{\frac{1}{\lambda}}-1}(1+t)^{\frac{x}{\lambda}}\frac{\mathrm{Li}_k\left(1-e^{-t}\right)}{t}.$$

We observe that

$$\begin{aligned}\frac{1}{t}\mathrm{Li}_k\left(1-e^{-t}\right) &= \frac{1}{t}\sum_{n=1}^{\infty}\frac{1}{n^k}\left(1-e^{-t}\right)^n \\ &= \frac{1}{t}\sum_{n=1}^{\infty}\frac{(-1)^n}{n^k}n!\sum_{l=n}^{\infty}S_2(l,n)\frac{(-t)^l}{l!} \\ &= \frac{1}{t}\sum_{l=1}^{\infty}\sum_{n=1}^{l}\frac{(-1)^{n+l}}{n^k}n!S_2(l,n)\frac{t^l}{l!} \\ &= \sum_{l=0}^{\infty}\sum_{n=1}^{l+1}\frac{(-1)^{n+l+1}}{n^k}n!\frac{S_2(l+1,n)}{l+1}\frac{t^l}{l!}.\end{aligned}\tag{2.7}$$

From (2.6) and (2.7), we have

$$\begin{aligned}&\sum_{n=0}^{\infty}\beta_n^{(k)}(x\mid\lambda)\frac{t^n}{n!} \\ &= \left(\sum_{m=0}^{\infty}\beta_m(x\mid\lambda)\frac{t^m}{m!}\right)\left(\sum_{l=0}^{\infty}\left(\sum_{p=1}^{l+1}\frac{(-1)^{p+l+1}}{p^k}p!\frac{S_2(l+1,p)}{l+1}\right)\frac{t^l}{l!}\right) \\ &= \sum_{n=0}^{\infty}\left\{\sum_{l=0}^{n}\binom{n}{l}\left(\sum_{p=1}^{l+1}\frac{(-1)^{p+l+1}p!}{p^k}\frac{S_2(l+1,p)}{l+1}\right)\beta_{n-l}(x\mid\lambda)\right\}\frac{t^n}{n!}.\end{aligned}\tag{2.8}$$

By comparing the coefficients on both sides of (2.8), we obtain the following theorem.

Theorem 2.2. *For $n \geq 0$, we have*

$$\beta_n^{(k)}(x\mid\lambda) = \sum_{l=0}^{n}\binom{n}{l}\left(\sum_{p=1}^{l+1}\frac{(-1)^{p+l+1}p!}{p^k}\frac{S_2(l+1,p)}{l+1}\right)\beta_{n-l}(x\mid\lambda).$$

It is easy to show that

$$\begin{aligned}&\frac{\mathrm{Li}_k\left(1-e^{-t}\right)}{(1+\lambda t)^{\frac{1}{\lambda}}-1}(1+\lambda t)^{\frac{x+1}{\lambda}} - \frac{\mathrm{Li}_k\left(1-e^{-t}\right)}{(1+\lambda t)^{\frac{1}{\lambda}}-1}(1+\lambda t)^{\frac{x}{\lambda}} \\ &= (1+\lambda t)^{\frac{x}{\lambda}}\mathrm{Li}_k\left(1-e^{-t}\right) \\ &= \left(\sum_{l=0}^{\infty}(x\mid\lambda)_l\frac{t^l}{l!}\right)\left(\sum_{m=1}^{\infty}\frac{\left(1-e^{-t}\right)^m}{m^k}\right) \\ &= \left(\sum_{l=0}^{\infty}(x\mid\lambda)_l\frac{t^l}{l!}\right)\left(\sum_{m=0}^{\infty}\frac{\left(1-e^{-t}\right)^{m+1}}{(m+1)^k}\right) \\ &= \left(\sum_{l=0}^{\infty}(x\mid\lambda)_l\frac{t^l}{l!}\right)\left(\sum_{p=1}^{\infty}\left(\sum_{m=0}^{p-1}\frac{(-1)^{m+p+1}}{(m+1)^k}(m+1)!S_2(p,m+1)\right)\frac{t^p}{p!}\right)\end{aligned}\tag{2.9}$$

$$= \sum_{n=1}^{\infty} \left\{ \sum_{p=1}^{n} \sum_{m=0}^{p-1} \frac{(-1)^{m+p+1}}{(m+1)^k} (m+1)! S_2 (p, m+1) \binom{n}{p} (x \mid \lambda)_{n-p} \right\} \frac{t^n}{n!}.$$

On the other hand,

$$\begin{aligned} &\frac{\mathrm{Li}_k \left(1-e^{-t}\right)}{(1+\lambda t)^{\frac{1}{\lambda}}-1} (1+\lambda t)^{\frac{x+1}{\lambda}} - \frac{\mathrm{Li}_k \left(1-e^{-t}\right)}{(1+\lambda t)^{\frac{1}{\lambda}}-1} (1+\lambda t)^{\frac{x}{\lambda}} \\ &= \sum_{n=0}^{\infty} \left\{ \beta_n^{(k)} (x+1 \mid \lambda) - \beta_n^{(k)} (x \mid \lambda) \right\} \frac{t^n}{n!}. \end{aligned} \tag{2.10}$$

Therefore, by (2.9) and (2.10), we obtain the following theorem.

Theorem 2.3. *For $n \geq 1$, we have*

$$\begin{aligned} &\beta_n^{(k)} (x+1 \mid \lambda) - \beta_n^{(k)} (x \mid \lambda) \\ &= \sum_{p=1}^{n} \left(\sum_{m=0}^{p-1} \frac{(-1)^{m+k+1}}{(m+1)^k} (m+1)! S_2 (k+m+1) \right) \binom{n}{p} (x \mid \lambda)_{n-p}. \end{aligned}$$

Now, we note that

$$\begin{aligned} &\frac{\mathrm{Li}_k \left(1-e^{-t}\right)}{(1+\lambda t)^{\frac{1}{\lambda}}-1} (1+\lambda t)^{\frac{x}{\lambda}} \\ &= \frac{\mathrm{Li}_k \left(1-e^{-t}\right)}{(1+\lambda t)^{\frac{d}{\lambda}}-1} \sum_{a=0}^{d-1} (1+\lambda t)^{\frac{l+x}{\lambda}} \\ &= \left(\frac{\mathrm{Li}_k \left(1-e^{-t}\right)}{t} \right) \frac{1}{d} \sum_{a=0}^{d-1} \frac{dt}{(1+\lambda t)^{\frac{d}{\lambda}}-1} (1+\lambda t)^{\frac{l+x}{\lambda}} \\ &= \sum_{l=0}^{\infty} \left(\sum_{p=1}^{l+1} \frac{(-1)^{p+l+1}}{p^k} p! \frac{S_2 (l+1, p)}{l+1} \right) \frac{t^l}{l!} \\ &\quad \times \sum_{a=0}^{d-1} \sum_{m=0}^{\infty} \beta_m \left(\frac{l+x}{d} \middle| \frac{\lambda}{d} \right) d^{m-1} \frac{t^m}{m!} \\ &= \sum_{a=0}^{d-1} \left(\sum_{n=0}^{\infty} \left(\sum_{l=0}^{n} \sum_{p=1}^{l+1} \binom{n}{l} \frac{(-1)^{p+l+1}}{p^k} p! \frac{S_2 (l+1, p)}{l+1} \beta_{n-l} \left(\frac{l+x}{d} \middle| \frac{\lambda}{d} \right) d^{n-l-1} \right) \frac{t^n}{n!} \right) \\ &= \sum_{n=0}^{\infty} \left\{ \sum_{a=0}^{d-1} \sum_{l=0}^{n} \sum_{p=1}^{l+1} \binom{n}{l} \frac{(-1)^{p+l+1}}{p^k} p! \frac{S_2 (l+1, p)}{l+1} \beta_{n-l} \left(\frac{l+x}{d} \middle| \frac{\lambda}{d} \right) d^{n-l-1} \right\} \frac{t^n}{n!}, \end{aligned} \tag{2.11}$$

where d is a fixed positive integer.

On the other hand,

$$\begin{aligned} &\frac{\mathrm{Li}_k \left(1-e^{-t}\right)}{(1+\lambda t)^{\frac{1}{\lambda}}-1} (1+\lambda t)^{\frac{x}{\lambda}} \\ &= \sum_{n=0}^{\infty} \beta_n^{(k)} (x \mid \lambda) \frac{t^n}{n!}. \end{aligned} \tag{2.12}$$

Therefore, by (2.11) and (2.12), we obtain the following theorem.

Theorem 2.4. *For $n \geq 0$, $d \in \mathbb{N}$ and $k \in \mathbb{Z}$, we have*

$$\begin{aligned}&\beta_n^{(k)}(x \mid \lambda)\\ &= \sum_{a=0}^{d-1}\sum_{l=0}^{n}\sum_{p=1}^{l+1}\binom{n}{l}\frac{(-1)^{p+l+1}}{p^k}p!\frac{S_2(l+1,p)}{l+1}\beta_{n-l}\left(\frac{l+x}{d}\middle|\frac{\lambda}{d}\right)d^{n-l-1}.\end{aligned}$$

From (2.4), we can derive the following equation:

$$\begin{aligned}&\sum_{n=0}^{\infty}\beta_n^{(k)}(x+y \mid \lambda)\frac{t^n}{n!}\\ &= \frac{\mathrm{Li}_k(1-e^{-t})}{(1+\lambda t)^{\frac{1}{\lambda}}-1}(1+\lambda t)^{\frac{x+y}{\lambda}}\\ &= \left(\frac{\mathrm{Li}_k(1-e^{-t})}{(1+\lambda t)^{\frac{1}{\lambda}}-1}(1+t\lambda)^{\frac{x}{\lambda}}\right)(1+\lambda t)^{\frac{y}{\lambda}}\\ &= \left(\sum_{l=0}^{\infty}\beta_l^{(k)}(x \mid \lambda)\frac{t^l}{l!}\right)\left(\sum_{m=0}^{\infty}(y \mid \lambda)_m\frac{t^m}{m!}\right)\\ &= \sum_{n=0}^{\infty}\left(\sum_{l=0}^{n}\binom{n}{l}\beta_l^{(k)}(x \mid \lambda)(y \mid \lambda)_{n-l}\right)\frac{t^n}{n!}.\end{aligned} \tag{2.13}$$

Therefore, by (2.13), we obtain the following theorem.

Theorem 2.5. *For $n \geq 0$, we have*

$$\beta_n^{(k)}(x+y \mid \lambda) = \sum_{l=0}^{n}\binom{n}{l}\beta_l^{(k)}(x \mid \lambda)(y \mid \lambda)_{n-l}.$$

Remark.

$$\begin{aligned}&\frac{d}{dx}\beta_n^{(k)}(x \mid \lambda)\\ &= \frac{d}{dx}\sum_{l=0}^{n}\binom{n}{l}\beta_{n-l}^{(k)}(\lambda)(x \mid \lambda)_l\\ &= \sum_{l=0}^{n}\binom{n}{l}\beta_{n-l}^{(k)}(\lambda)\sum_{j=0}^{l-1}\frac{1}{x-\lambda j}\prod_{i=0}^{l-1}(x-\lambda i)\\ &= \sum_{l=0}^{n}\binom{n}{l}\beta_{n-l}^{(k)}(\lambda)\sum_{j=0}^{l-1}\prod_{\substack{i=0\\ i\neq j}}^{l-1}(x-\lambda i).\end{aligned}$$

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to this work. All authors read and approved the final manuscript.

References

1. M.t Açikgöz, D. Erdal, and S. Araci, *A new approach to q-Bernoulli numbers and q-Bernoulli polynomials related to q-Bernstein polynomials*, Adv. Difference Equ. (2010), Art. ID 951764, 9. MR 2764538 (2012g:11038)
2. S. Araci, M. Acikgoz, and H. Jolany, *On the families of q-Euler polynomials and their applications*, J. Egyptian Math. Soc. **23** (2015), no. 1, 1–5. MR 3317290
3. A. Bayad and S. Gaboury, *Generalized Dirichlet L-function of arbitrary order with applications*, Adv. Stud. Contemp. Math. (Kyungshang) **23** (2013), no. 4, 607–619. MR 3115219
4. A. Bayad and Y. Hamahata, *Multiple polylogarithms and multi-poly-Bernoulli polynomials*, Funct. Approx. Comment. Math. **46** (2012), no. part 1, 45–61. MR 2951728
5. L. Carlitz, *A degenerate Staudt-Clausen theorem*, Arch. Math. (Basel) **7** (1956), 28–33. MR 0074436 (17,586a)
6. ______, *Degenerate Stirling, Bernoulli and Eulerian numbers*, Utilitas Math. **15** (1979), 51–88. MR 531621 (80i:05014)
7. R. Dere and Y. Simsek, *Applications of umbral algebra to some special polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **22** (2012), no. 3, 433–438. MR 2976601
8. D. Ding and J. Yang, *Some identities related to the Apostol-Euler and Apostol-Bernoulli polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **20** (2010), no. 1, 7–21. MR 2597988 (2011k:05030)
9. S. Gaboury, R. Tremblay, and B.-J. Fugère, *Some explicit formulas for certain new classes of Bernoulli, Euler and Genocchi polynomials*, Proc. Jangjeon Math. Soc. **17** (2014), no. 1, 115–123. MR 3184467
10. H. Jolany and H. Mohebbi, *Some results on Generalized multi poly-Bernoulli and Euler polynomials*, Math. Combin. Book Ser. **2** (2011), 117–129.
11. D. S. Kim and T. Kim, *Hermite and poly-Bernoulli mixed-type polynomials*, Adv. Difference Equ. (2013), 2013:343, 12. MR 3213908
12. ______, *Higher-order Frobenius-Euler and poly-Bernoulli mixed-type polynomials*, Adv. Difference Equ. (2013), 2013:251. MR 3108262
13. ______, *A note on poly-Bernoulli and higher-order poly-Bernoulli polynomials*, Russ. J. Math. Phys. **22** (2015), no. 1, 26–33. MR 3318873
14. ______, *Some identities of degenerate Euler polynomials arising from p-adic fermionic integrals on $\mathbb{Z}_p$*, Integral Transforms Spec. Funct. **26** (2015), no. 4, 295–302. MR 3304355
15. D. S. Kim, T. Kim, D. V. Dolgy, and T. Komatsu, *Barnes-type degenerate Bernoulli polynomials*, Adv. Stud. Contemp. Math. **25** (2015), no. 1, 121–146.
16. D. S. Kim, T. Kim, and S. H. Lee, *A note on poly-bernoulli polynomials arising umbral calculus*, Adv. Studies Theor. Phys. **7** (2013), no. 15, 731–744.
17. D. S. Kim, T. Kim, T. Mansour, and D. V. Dolgy, *On Poly-Bernoulli polynomials of the second kind with umbral calculus viewpoint*, Adv. Difference Equ. (2015), no. 1, 2015:27. MR 3304954

18. T. Kim, *Some identities on the q-Euler polynomials of higher order and q-Stirling numbers by the fermionic p-adic integral on* $\mathbb{Z}_p$, Russ. J. Math. Phys. **16** (2009), no. 4, 484–491. MR 2587805 (2011e:33045)
19. ______, *Barnes' type multiple degenerate Bernoulli and Euler polynomials*, Applied Math. Comput. **258** (2015), 556–564.
20. T. Kim, Y. S. Jang, and J. J. Seo, *poly-Bernoulli polynomials and their applications*, Int. Journal of Math. Analysis **8** (2014), no. 30, 1495–1503.
21. T. Kim, H. K. Kwon, S. H. Lee, and J. J. Seo, *A note on poly-Bernoulli numbers and polynomials of the second kind*, Adv. Difference Equ. (2014), 2014:219.
22. Q.-M. Luo, B.-N. Guo, and F. Qi, *On evaluation of Riemann zeta function* $\zeta(s)$, Adv. Stud. Contemp. Math. (Kyungshang) **7** (2003), no. 2, 135–144. MR 2005861 (2004h:11069)
23. Q.-M. Luo and F. Qi, *Relationships between generalized Bernoulli numbers and polynomials and generalized Euler numbers and polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **7** (2003), no. 1, 11–18. MR 1981601
24. J.-W. Park and S.-H. Rim, *On the modified q-Bernoulli polynomials with weight*, Proc. Jangjeon Math. Soc. **17** (2014), no. 2, 231–236. MR 3234991
25. E. Şen, *Theorems on Apostol-Euler polynomials of higher order arising from Euler basis*, Adv. Stud. Contemp. Math. (Kyungshang) **23** (2013), no. 2, 337–345. MR 3088764
26. H. M. Srivastava, T. Kim, and Y. Simsek, *q-Bernoulli numbers and polynomials associated with multiple q-zeta functions and basic L-series*, Russ. J. Math. Phys. **12** (2005), no. 2, 241–268. MR 2199005 (2006k:11034)
27. Z. Zhang and J. Yang, *On sums of products of the degenerate Bernoulli numbers*, Integral Transforms Spec. Funct. **20** (2009), no. 9-10, 751–755.

Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea
E-mail address: `dskim@sogang.ac.kr`

Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea
E-mail address: `tkkim@kw.ac.kr`